

JOURNAL OF  
**Neurobiology**

**Volume Contents**

*Vol. 51, No. 1, April 2002*

**Antiandrogen Blocks Estrogen-Induced Masculinization of the Song System in Female Zebra Finches / 1**

*William Grisham, Janet Lee, Mary Ellen McCormick, Kay Yang-Stayner, and Arthur P. Arnold*

**Long-Term Culture of Mouse Cortical Neurons as a Model for Neuronal Development, Aging, and Death / 9**

*Christian Lesuisse and Lee J. Martin*

**Progesterone Receptors and the Sexual Differentiation of the Medial Preoptic Nucleus / 24**

*Princy S. Quadros, Veronica Lopez, Geert J. De Vries, Wilson C. J. Chung, and Christine K. Wagner*

**Perineal Muscles and Motoneurons Are Sexually Monomorphic in the Naked Mole-Rat (*Heterocephalus glaber*) / 33**

*Maria E. Peroulakis, Bruce Goldman, and Nancy G. Forger*

**Temporal Regulation of Neuropilin-1 Expression and Sensitivity to Semaphorin 3A in NGF- and NT3-Responsive Chick Sensory Neurons / 43**

*Ausra Pond, Florence K. Roche, and Paul C. Letourneau*

**Regulated Appearance of NMDA Receptor Subunits and Channel Functions during *In Vitro* Neuronal Differentiation / 54**

*Márta Jelítai, Katalin Schlett, Patrícia Varju, Ulrich Eisel, and Emília Madarász*

**Inositol 1,4,5-Trisphosphate Transduction Cascade in Taste Reception of the Fleshfly, *Boettcherisca peregrina* / 66**

*Masayuki Koganezawa and Ichiro Shimada*

**Cover:**  $\beta$ -synuclein is present in dendritic growth cones in developing cortical neurons. Mouse cortical neuron cultures at 3 days in vitro were used to immunolocalize AMPA receptor GluR1 (red staining) and  $\beta$ -synuclein (green staining). In dendritic growth cones,  $\beta$ -synuclein is discretely localized at the base of the growth cone (see dendritic growth cones at left of cell detected with GluR1 antibodies). This localization provides insight into the possible functions of  $\beta$ -synuclein. See article by Lesuisse and Martin on page 9.

**Vol. 51, No. 2, May 2002**

**FAST TRACK**

**Stimulation of Nitric Oxide Synthase in Cerebral Cortex Due to Elevated Partial Pressures of Oxygen: An Oxidative Stress Response / 85**

*Stephen R. Thom, Veena Bhopale, Donald Fisher, Yefim Manevich, Paul L. Huang, and Donald G. Buerk*

**Comparison of the Effects of HGF, BDNF, CT-1, CNTF, and the Branchial Arches on the Growth of Embryonic Cranial Motor Neurons / 101**

*Arifa Naeem, Leila Abbas, and Sarah Guthrie*

**The Effects of Social Environment on Adult Neurogenesis in the Female Prairie Vole / 115**

*Christie D. Fowler, Yan Liu, Charles Ouimet, and Zuoxin Wang*

**Respiratory Units of Motor Production and Song Imitation in the Zebra Finch / 129**

*Michele Franz and Franz Goller*

**Changes in Spatial Memory Mediated by Experimental Variation in Food Supply Do Not Affect Hippocampal Anatomy in Mountain Chickadees (*Poecile gambeli*) / 142**

*V. V. Pravosudov, P. Lavenex, and N. S. Clayton*

**Developmental and Hormonal Regulation of NR2A mRNA in Forebrain Regions Controlling Avian Vocal Learning / 149**

*J. E. Heinrich, T. D. Singh, F. Sohrabji, K. W. Nordeen, and E. J. Nordeen*

**Influence of Ethanol on Neonatal Cerebellum of BDNF Gene-Deleted Animals: Analyses of Effects on Purkinje Cells, Apoptosis-Related Proteins, and Endogenous Antioxidants / 160**

*Marieta Barrow Heaton, Irina Madorsky, Michael Paiva, and Joanne Mayer*

**Cover:** A confocal laser microscope image of cells colocalized (yellow fluorescence) for a cell proliferation marker, BrdU (red fluorescence), and a neuronal marker, TuJ1 (green fluorescence), in the subventricular zone of a female prairie vole. From the article by Fowler et al. on page 115.

**Vol. 51, No. 3, June 5, 2002**

**FAST TRACK**

**L1 Mediated Homophilic Binding and Neurite Outgrowth Are Modulated by Alternative Splicing Exon 2 / 177**

*Jeffrey Jacob, Jeffrey Haspel, Noriko Kane-Goldsmith, and Martin Grumet*

**Complementary Expression and Heterophilic Interactions between IgLON Family Members Neurotrimin and LAMP / 190**

*Orlando D. Gil, Li Zhang, Suzanne Chen, Y. Q. Ren, Aurea Pimenta, George Zanazzi, Dean Hillman, Pat Levitt, and James L. Salzer*

**Forked End: A Novel Transmembrane Protein Involved in Neuromuscular Specificity in *Drosophila* Identified by Gain-of-Function Screening / 205**

*Takeshi Umekiya, Etsuko Takasu, Masatoshi Takeichi, Toshiro Aigaki, and Akinao Nose*

**Septum Volume and Food-Storing Behavior Are Related in Parids / 215**

*Michael W. Shiftlett, Kristy L. Gould, Tom V. Smulders, and Timothy J. DeVoogd*

**Time Course of Morphological Alterations of Fungiform Papillae and Taste Buds Following Chorda Tympani Transection in Neonatal Rats / 223**

*Suzanne I. Sollars, Peter C. Smith, and David L. Hill*

**Multiphoton Stimulation of Neurons / 237**

*Hajime Hirase, Volodymyr Nikolenko, Jesse H. Goldberg, and Rafael Yuste*

**Inactivation of Olfactory Sensilla of a Single Morphological Type Differentially Affects the Response of *Drosophila* to Odors / 248**

*S.-K. Park, S. R. Shanbhag, A. E. Dubin, M. de Bruyne, Q. Wang, P. Yu, N. Shimoni, S. D'Mello, J. R. Carlson, G. L. Harris, R. A. Steinbrecht, and C. W. Pikielny*

**Cover:** Coaggregation of beads coated with proteins shows that L1 beads (red) bind well to themselves forming large aggregates but only weakly to beads coated with a splicing variant lacking exon 2 called  $\Delta 2$  (green);  $\Delta 2$  is associated with a human mutation in L1. From the article by Jacob et al. on pages 177–189.

**Vol. 51, No. 4, June 15, 2002**

**FAST TRACK**

**Dominant-Negative NSF2 Disrupts the Structure and Function of *Drosophila* Neuromuscular Synapses / 261**

*Bryan A. Stewart, Mahmood Mohtashami, Patricia Rivlin, David L. Deitcher, William S. Trimble, and Gabrielle L. Boulianne*

**GABA and Development of the *Xenopus* Optic Projection / 272**

*Shane C. D. Ferguson and Sarah McFarlane*

**Binding Characteristics of Chondroitin Sulfate Proteoglycans and Laminin-1, and Correlative Neurite Outgrowth Behaviors in a Standard Tissue Culture Choice Assay / 285**

*Diane M. Snow, Jeffrey D. Smith, and Julie A. Gurwell*

**High-Resolution Imaging Demonstrates Dynein-Based Vesicular Transport of Activated Trk Receptors / 302**

*Anita Bhattacharyya, Fiona L. Watson, Scott L. Pomeroy, Yanzen Z. Zhang, Charles D. Stiles, and Rosalind A. Segal*

**Decreased Calretinin Expression in Cerebellar Granule Cells in the Leaner Mouse / 313**

*Sang-Soep Nahm, Dana J. Tomlinson, and Louise C. Abbott*

**Effects of Excess Thyroid Hormone on Cell Death, Cell Proliferation, and New Neuron Incorporation in the Adult Zebra Finch Telencephalon / 323**

*P. K. Tekumalla, M. Tontonoz, M. A. Hesla, and J. R. Kirn*

**Author Index to Volume 51 / 345**

**Subject Index to Volume 51 / 347**

**Volume Contents / I**

**Cover:** High resolution imaging demonstrates colocalization of dynein components and activated Trk receptors in axons. Longitudinal sections of rat sciatic nerve were immunostained for activated Trk (in red) and dynamin, a component of the dynactin complex (in green). Deconvolved image of a single optical section shows dynamin within the Schwann cells (green), surrounding individual axons containing punctate phospho-Trk (red). Within the axons, many phospho-Trk containing vesicles colocalize with dynein and dynamin (yellow). See article by Bhattacharyya et al., p. 302.

